



**blood**

2014 124: 1404-1411

doi:10.1182/blood-2014-03-565135 originally published  
online July 15, 2014

## **Treatment recommendations for patients with Waldenström macroglobulinemia (WM) and related disorders: IWWM-7 consensus**

Meletios A. Dimopoulos, Efsthios Kastiris, Roger G. Owen, Robert A. Kyle, Ola Landgren, Enrica Morra, Xavier Leleu, Ramón García-Sanz, Nikhil Munshi, Kenneth C. Anderson, Evangelos Terpos, Irene M. Ghobrial, Pierre Morel, David Maloney, Mathias Rummel, Véronique Leblond, Ranjana H. Advani, Morie A. Gertz, Charalampia Kyriakou, Sheeba K. Thomas, Bart Barlogie, Stephanie A. Gregory, Eva Kimby, Giampaolo Merlini and Steven P. Treon

---

Updated information and services can be found at:

<http://www.bloodjournal.org/content/124/9/1404.full.html>

Articles on similar topics can be found in the following Blood collections

[Free Research Articles](#) (2720 articles)

[Lymphoid Neoplasia](#) (1827 articles)

[Multiple Myeloma](#) (247 articles)

[Review Articles](#) (529 articles)

---

Information about reproducing this article in parts or in its entirety may be found online at:

[http://www.bloodjournal.org/site/misc/rights.xhtml#repub\\_requests](http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests)

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>

## Review Article

# Treatment recommendations for patients with Waldenström macroglobulinemia (WM) and related disorders: IWWM-7 consensus

Meletios A. Dimopoulos,<sup>1</sup> Efstathios Kastritis,<sup>1</sup> Roger G. Owen,<sup>2</sup> Robert A. Kyle,<sup>3</sup> Ola Landgren,<sup>4</sup> Enrica Morra,<sup>5</sup> Xavier Leleu,<sup>6</sup> Ramón García-Sanz,<sup>7</sup> Nikhil Munshi,<sup>8</sup> Kenneth C. Anderson,<sup>8</sup> Evangelos Terpos,<sup>1</sup> Irene M. Ghobrial,<sup>8</sup> Pierre Morel,<sup>9</sup> David Maloney,<sup>10</sup> Mathias Rummel,<sup>11</sup> Véronique Leblond,<sup>12</sup> Ranjana H. Advani,<sup>13</sup> Morie A. Gertz,<sup>3</sup> Charalampia Kyriakou,<sup>14</sup> Sheeba K. Thomas,<sup>15</sup> Bart Barlogie,<sup>16</sup> Stephanie A. Gregory,<sup>17</sup> Eva Kimby,<sup>18</sup> Giampaolo Merlini,<sup>19</sup> and Steven P. Treon<sup>8</sup>

<sup>1</sup>Department of Clinical Therapeutics, National and Kapodistrian University of Athens School of Medicine, Athens, Greece; <sup>2</sup>St. James's Institute of Oncology, Leeds, United Kingdom; <sup>3</sup>Mayo Clinic, Rochester, MN; <sup>4</sup>National Cancer Institute, Bethesda, MD; <sup>5</sup>Department of Hematology, Niguarda Ca' Granda Hospital, Milan, Italy; <sup>6</sup>Service des Maladies du Sang, Hôpital Huriez, Centre Hospitalier Régional Universitaire, Lille, France; <sup>7</sup>Servicio de Hematología, Hospital Universitario de Salamanca, Instituto de Investigación Biomédica de Salamanca, Salamanca, Spain; <sup>8</sup>Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA; <sup>9</sup>Service d'Hématologie Clinique, Centre Hospitalier Schaffner, Lens, France; <sup>10</sup>Fred Hutchinson Cancer Research Center, Seattle, WA; <sup>11</sup>Medizinischen Klinik IV, Hospital of the Justus-Liebig-University Giessen, Germany; <sup>12</sup>Assistance Publique-Hôpitaux de Paris Hôpital Pitié-Salpêtrière, Université Pierre et Marie Curie Paris 6 GRC-11, Groupe de Recherche Clinique Hémopathie Lymphoïde, Paris, France; <sup>13</sup>Stanford Cancer Institute, Stanford, CA; <sup>14</sup>Royal Free Hospital, London, United Kingdom; <sup>15</sup>University of Texas MD Anderson Cancer Center, TX; <sup>16</sup>Myeloma Institute for Research and Therapy, University of Arkansas for Medical Sciences, Little Rock, AR; <sup>17</sup>Rush University Medical Center, Chicago, IL; <sup>18</sup>Department of Hematology, Karolinska University Hospital, Solna, Sweden; and <sup>19</sup>Amyloidosis Research and Treatment Center, Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Policlinico San Matteo, Department of Molecular Medicine, University of Pavia, Pavia, Italy

**Waldenström macroglobulinemia (WM) is a distinct B-cell lymphoproliferative disorder for which clearly defined criteria for the diagnosis, initiation of therapy, and treatment strategy have been proposed as part of the consensus panels of International Workshops on WM (IWWM). As part of the IWWM-7 and based on recently published and ongoing clinical trials, the panels updated treatment recommendations. Therapeutic strategy in WM should be based on individual patient and disease characteristics (age, comorbidities, need for rapid disease control, candidacy for**

**autologous transplantation, cytopenias, IgM-related complications, hyperviscosity, and neuropathy). Mature data show that rituximab combinations with cyclophosphamide/dexamethasone, bendamustine, or bortezomib/dexamethasone provided durable responses and are indicated for most patients. New monoclonal antibodies (ofatumumab), second-generation proteasome inhibitors (carfilzomib), mammalian target of rapamycin inhibitors, and Bruton's tyrosine kinase inhibitors are promising and may expand future treatment options. A different regimen is**

**typically recommended for relapsed or refractory disease. In selected patients with relapsed disease after long-lasting remission, reuse of a prior effective regimen may be appropriate. Autologous stem cell transplantation may be considered in young patients with chemosensitive disease and in newly diagnosed patients with very-high-risk features. Active enrollment of patients with WM in clinical trials is encouraged. (Blood. 2014;124(9):1404-1411)**

## Introduction

Waldenström macroglobulinemia (WM) is, according to the World Health Organization classification, a lymphoplasmacytic lymphoma<sup>1</sup> in which the bone marrow is infiltrated by immunoglobulin (Ig)M-producing clonal lymphoplasmacytic cells. The Second International Workshop on WM (IWWM-2) proposed criteria for the clinicopathological diagnosis and for initiation of therapy in WM patients.<sup>2,3</sup> The IWWM consensus panels have provided treatment recommendations,<sup>4,5</sup> which were last updated in 2008 (IWWM-4).<sup>6</sup> As part of its last consensus deliberations (IWWM-7, Newport, RI, August 2012), the panel considered the results from phase 2 studies of several chemoimmunotherapy regimens, novel drugs (alone or with rituximab), and emerging novel targeted agents (ofatumumab, everolimus, perifosine, enzastaurin, panobinostat, carfilzomib, and ibrutinib); examined these data; and updated its recommendations, which are presented herein.

The consensus panels recommended that individual patient considerations should be weighed for the choice of therapy, including

the need for rapid disease control, age, candidacy for autologous transplantation, comorbidities, presence of cytopenias, hyperviscosity, lymphadenopathy, IgM-related end-organ damage, and patients' preferences. Based on available data, the panel provides guidance on the management of patients with WM adjusted to specific conditions and complications of the disease both for the initial therapy and for relapsed or refractory disease.

### Major changes since the last published recommendations

Rituximab-based regimens remain a recommended primary therapy for most patients with WM. As per the previous recommendations of IWWM-4,<sup>6</sup> dexamethasone, rituximab, and cyclophosphamide (DRC) remains a primary choice, but combinations such as rituximab-cyclophosphamide, doxorubicin, vincristine, and prednisone (R-CHOP) are no longer considered a first-line choice; instead,

**Table 1. Indications for initiation of therapy in patients with WM**

<b>Clinical indications for initiation of therapy</b>	
Recurrent fever, night sweats, weight loss, fatigue	
Hyperviscosity	
Lymphadenopathy which is either symptomatic or bulky ( $\geq 5$ cm in maximum diameter)	
Symptomatic hepatomegaly and/or splenomegaly	
Symptomatic organomegaly and/or organ or tissue infiltration	
Peripheral neuropathy due to WM	
<b>Laboratory indications for initiation of therapy</b>	
Symptomatic cryoglobulinemia	
Cold agglutinin anemia	
Immune hemolytic anemia and/or thrombocytopenia	
Nephropathy related to WM	
Amyloidosis related to WM	
Hemoglobin $\leq 10$ g/dL	
Platelet count $< 100 \times 10^9/L$	

bendamustine-rituximab (BR) is now a primary treatment option, especially for patients with high tumor bulk. In the current recommendations bortezomib-rituximab combinations may also be considered a primary option for patients with specific high-risk features (ie, hyperviscosity) or in younger patients for whom avoidance of alkylator therapy is sought. Fludarabine-based combinations are not recommended for primary therapy but remain an option for patients with relapsed/refractory disease with adequate performance status. In patients who may be candidates for single agent oral therapy, oral fludarabine (if available) is recommended over chlorambucil.

### Risk stratification

The importance of a prognostic system for the risk stratification of patients with WM and as a tool for study comparisons has been emphasized.<sup>6</sup> In International Prognostic Scoring System for WM I (IPSSWM), 5 covariates (age  $> 65$  years, hemoglobin  $\leq 11.5$  g/dL, platelet counts  $\leq 100 \times 10^9/L$ ,  $\beta 2$ -microglobulin  $> 3$  mg/L, serum monoclonal protein  $> 70$  g/L) defined 3 risk groups (low, intermediate, and high risk, respectively).<sup>7</sup> IPSSWM has been validated externally, and its prognostic significance has been confirmed.<sup>8-10</sup> Results per IPSSWM risk category are increasingly reported and are used for stratification in randomized clinical trials. However, the use of IPSSWM in making treatment decisions remains to be delineated.

### Justifying treatment initiation

Not all patients with a diagnosis of WM need immediate therapy. Criteria for the initiation of therapy (proposed in the IWWM-2 consensus panel and confirmed in IWWM-7) are presented in Table 1. For patients who do not fulfill the criteria in Table 1 and in whom only laboratory evidence may indicate a possible development of symptomatic disease (such as a minor decrease in hemoglobin level, but  $> 10$  g/dL, or mild increases in IgM or mild increase of lymphadenopathy or splenomegaly without discomfort for the patient), close observation is recommended.<sup>3</sup>

### Risk assessment for progression to symptomatic disease and follow-up recommendations

IgM-monoclonal gammopathy of undetermined significance or asymptomatic WM are increasingly diagnosed because more individuals undergo a serum protein electrophoresis as part of a routine laboratory assessment. The diagnosis of asymptomatic WM requires the demonstration of infiltration of the bone marrow by  $\geq 10\%$

clonal lymphoplasmacytic cells on trephine biopsy or a monoclonal IgM  $> 3$  g/dL and no end-organ damage or symptoms.<sup>11</sup>

The median time to initiation of therapy for asymptomatic patients in the Southwest Oncology Group-S9003 study exceeded 7 years.<sup>12</sup> In the series by Kyle et al, the cumulative probability of progression for patients with asymptomatic WM was 6%, 39%, 59%, and 68% at 1, 3, 5, and 10 years, respectively,<sup>11</sup> and 75% required therapy during a median of 15 years of follow-up. Lower hemoglobin,<sup>11,12</sup> extensive bone marrow infiltration,<sup>11</sup> size of serum M-spike,<sup>11</sup> and  $\beta$ -microglobulin levels<sup>12</sup> were significant predictors of an eventual need for therapy.

There are no data to justify early initiation of treatment, and patients with asymptomatic WM should be followed without therapy, preferably every 3 months for the first year to evaluate the pace of disease progression and, if stable, at more extended intervals thereafter. The risk of progression remains, and individuals with asymptomatic WM should be followed for life.

### Evaluation of response to therapy

The consensus-based uniform response criteria for WM, based mainly on the degree of M-protein reduction, were recently updated<sup>5,6,13</sup> (Table 2). Caution is advised in the early evaluation of response during rituximab-based therapy (or other anti-CD20 monoclonal antibodies), because of the common IgM flare,<sup>14-16</sup> which does not necessarily imply disease progression; in most cases, it will resolve, but if necessary, additional tests may be performed to discriminate from disease progression. In patients treated with agents such as bortezomib or everolimus, tumor reduction in the bone marrow may not be proportional to the suppression of IgM levels.<sup>17-20</sup> Thus, the variability of the IgM kinetics with various therapies should be taken into account, and in discordant cases, additional investigations should be considered.

## New treatment options for patients with WM

### Bendamustine

In the study group of indolent lymphomas/non-Hodgkin lymphoma (StiL NHL) 1-2003 study, 513 newly diagnosed patients with follicular, marginal zone, small lymphocytic, and mantle cell lymphoma and WM received BR or R-CHOP: of 41 patients with WM, 22 received BR and 19 received R-CHOP.<sup>21</sup> Responses were similar (95% in both arms), but BR was superior in terms of progression-free survival (PFS; median, 69.5 vs 28.1 months;  $P = .0033$ ) and tolerability. After a median follow-up of 45 months, a difference in overall survival was observed after the fifth year, but further follow-up is needed. Regarding stem cell harvest, after 6 cycles of BR, the CD34 yield was similar to that after R-CHOP.

As part of a large study investigating rituximab maintenance in patients with previously untreated low-grade lymphomas (including WM), BR induction was given in 162 patients with WM (116 were evaluable for response), and 86% achieved at least partial response (PR).<sup>22</sup> Responding patients (at least PR) were randomized to either observation or 2 years of rituximab maintenance; updated results are awaited.

Bendamustine is also active in patients with relapsed or refractory WM, either with rituximab, ofatumumab, or as monotherapy (very good partial response [VGPR] in 17%, PR in 67%; median PFS, 13.2 months), but prior nucleoside analog exposure was associated with prolonged myelosuppression.<sup>23</sup>

**Table 2. Consensus-based uniform response criteria for WM developed by the IWWM, updated in the sixth IWWM**

Response category	Description
CR	Absence of serum monoclonal IgM protein by immunofixation Normal serum IgM level Complete resolution of extramedullary disease, ie, lymphadenopathy and splenomegaly if present at baseline Morphologically normal bone marrow aspirate and trephine biopsy
VGPR	Monoclonal IgM protein is detectable ≥90% reduction in serum IgM level from baseline* Complete resolution of extramedullary disease, ie, lymphadenopathy/splenomegaly if present at baseline No new signs or symptoms of active disease
PR	Monoclonal IgM protein is detectable ≥50% but <90% reduction in serum IgM level from baseline* Reduction in extramedullary disease, ie, lymphadenopathy/splenomegaly if present at baseline No new signs or symptoms of active disease
MR	Monoclonal IgM protein is detectable ≥25% but <50% reduction in serum IgM level from baseline* No new signs or symptoms of active disease
Stable disease	Monoclonal IgM protein is detectable <25% reduction and <25% increase in serum IgM level from baseline* No progression in extramedullary disease, ie, lymphadenopathy/splenomegaly No new signs or symptoms of active disease
Progressive disease	≥25% increase in serum IgM level* †from lowest nadir (requires confirmation) and/or progression in clinical features attributable the disease

Published by Owen et al.<sup>13</sup>

\*Sequential changes in IgM levels may be determined either by M-protein quantitation by densitometry or total serum IgM quantitation by nephelometry.

†An absolute increase of &gt;5 g/L (0.5 g/dL) is required when the increase of IgM component is the only applicable criterion.

Thus, available data indicate that BR is at least as effective as R-CHOP, may be associated with longer PFS and less toxicity, and probably does not compromise stem cell collection. However, R-CHOP is not a standard first-line regimen for WM<sup>24</sup>; less intensive and less toxic regimens like DRC are more often used. No increased incidence of secondary malignancies after BR was observed but longer follow-up is needed.<sup>21</sup> BR is a primary option for patients with newly diagnosed WM, especially those in need for rapid disease control or with bulky disease.<sup>25</sup>

### Bortezomib

Single agent bortezomib is associated with responses in ~40% of patients with relapsed or refractory WM, including fludarabine and rituximab refractory patients. Bortezomib with rituximab (±dexamethasone) are active, and in the front-line setting, 3 phase 2 studies of bortezomib/rituximab combinations showed response rates (at least PR) between 66% and 83% and rapid times to first response (2-3 months). Treon et al<sup>26</sup> treated 23 previously untreated patients with 4 consecutive cycles of bortezomib, dexamethasone, and rituximab (BDR) using twice-weekly, intravenous bortezomib, followed by 4 additional cycles given at 3-month intervals (at least

PR in 83% and complete response [CR]/near CR [nCR] in 22%); 61% of patients discontinued bortezomib due to neurotoxicity, whereas 2 patients developed symptomatic IgM flare. To reduce neurotoxicity, Ghobrial et al used weekly bortezomib with rituximab (6 cycles, no maintenance) in 26 patients<sup>27</sup> (58% achieved at least PR, 8% CR/nCR, and 1-year event-free survival was 79%). IgM flare developed in 40% and 54% of patients who developed peripheral neuropathy, but in none was the grade ≥3. Dimopoulos et al, to avoid IgM flare, used an induction cycle of bortezomib (intravenous 1.3 mg/m<sup>2</sup> on days 1, 4, 8, and 11), followed by 4 cycles of weekly bortezomib (intravenous 1.6 mg/m<sup>2</sup> for 4 weeks) with rituximab and dexamethasone on cycles 2 and 5. Among 59 previously untreated patients, 68% achieved at least PR (3% CR, 7% VGPR); IgM flare occurred in 11%, but plasmapheresis was not required, probably due to the initial bortezomib induction. After a median follow-up of 42 months, responses were durable (median PFS was 42 months and 3-year PFS for those with at least PR was 70%), despite the lack of maintenance. Peripheral neuropathy was observed in 46% (grade ≥ 3 in 7%), but only 5 (8%) patients discontinued bortezomib due to neuropathy.

Neurotoxicity is the major concern with bortezomib because underlying IgM-related neuropathy or neuropathies due to age-related comorbidities (such as diabetes) are common. Weekly dosing<sup>28-30</sup> and subcutaneous administration may reduce rates and severity of neuropathy and is explored in a clinical trial (NCT01592981). Bortezomib is not stem cell toxic, and long-term follow-up in myeloma patients does not suggest a risk for secondary malignancies.<sup>31</sup> Prophylaxis against herpes zoster is strongly recommended. Primary therapy with bortezomib is recommended for patients with high levels of IgM, with symptoms of, or at risk of developing, hyperviscosity syndrome, symptomatic cryoglobulinemia or cold agglutininemia, amyloidosis, and renal impairment.

Carfilzomib, a second-generation proteasome inhibitor, is associated with a low risk of neurotoxicity in myeloma patients and was recently evaluated in combination with rituximab and dexamethasone (CaRD), mainly in untreated WM patients.<sup>32</sup> The schedule of carfilzomib was attenuated (days 1, 2, 8, and 9) compared with myeloma dosing, and maintenance therapy (days 1 and 2 only) was given every 8 weeks for 8 cycles. The overall response rate was 87% (at least VGPR in 35%), and no grade ≥3 neuropathy was observed. With a median follow-up of 15.4 months, 20 of 31 (65%) patients remained progression free. CaRD therefore represents a novel neuropathy-sparing option for proteasome inhibitor-based therapy for WM.

### Everolimus

Long-term results from a phase 2 study of the oral mammalian target of rapamycin (mTOR) inhibitor everolimus (10 mg daily) showed at least PR in 50% (additional 23% minor responses [MRs]) of previously treated patients. Responses were rapid, occurring at a median of 2 months, and the median PFS was 21 months.<sup>33</sup> Grade ≥ 3 toxicities occurred in 67% of patients. Among previously untreated patients (N = 33), everolimus induced at least PR in 61% (plus 12% MRs),<sup>20</sup> although discordance between serum IgM levels and bone marrow disease burden was commonly observed. Six (18%) patients discontinued therapy due to toxicity. Myelosuppression with everolimus was common, and significant nonhematological toxicities included diarrhea, fatigue, and stomatitis (8-27%). Additionally, 5% to 15% developed pulmonary toxicity, frequently resulting in interruption or discontinuation of therapy. Everolimus

**Table 3. Summary of changes from the previous recommendations from the fourth IWWM, published in 2009**

Clinical condition	New recommendation (2014)	Old recommendation (2009)
Cytopenias	DRC, bendamustine-rituximab, bortezomib-rituximab	DRC, thalidomide + rituximab
High M-protein, transplant candidate	Bendamustine-rituximab, bortezomib-rituximab	R-CHOP, DRC
High M-protein, non-transplant candidate	Bendamustine-rituximab, bortezomib-rituximab	Nucleoside analogs + rituximab; nucleoside analogs + rituximab + cyclophosphamide
Comorbidities and cytopenias	Rituximab	Rituximab
Older age, slow progression, candidate for oral therapy	Oral fludarabine	Cladribine

may therefore be considered for selected patients with relapsed or refractory disease and limited options.

### Fludarabine

Nucleoside analogs have been used extensively in the treatment of WM, either alone or in different combinations. In the large WM1 randomized phase 3 study, oral fludarabine was superior to chlorambucil in terms of response rates, PFS, and overall survival (OS) in newly diagnosed WM/lymphoplasmacytic lymphoma.<sup>34</sup> Fludarabine has been studied in combination with rituximab, with or without cyclophosphamide (FR or FCR). Tedeschi et al<sup>35</sup> published long-term results (median follow-up, 37 months) with FCR in either previously untreated patients or patients relapsing after 1 line of alkylator-based (chlorambucil or cyclophosphamide) therapy. At least PR was observed in 74.4% (CR in 11.6% and VGPR in 20.9%), with an improvement of the quality of responses observed during follow-up, to a final 18.6% of CRs. Median event-free survival was 50 months, and estimated 4-year OS was 69%. Myelosuppression, especially neutropenia, was the main reason for treatment discontinuation, including episodes of long-lasting neutropenia after the end of treatment, whereas 3 patients developed myelodysplastic syndrome. In retrospective studies, fludarabine-based therapy was associated with an increased risk of secondary malignancies,<sup>36</sup> although the frequency was not greater than with chlorambucil in the WM1 study.<sup>34</sup> Nevertheless, response rates with fludarabine combinations are high, even in patients with relapsed or refractory WM, and the duration of response is long. Fludarabine-based combinations should be considered in patients with relapsed/refractory WM, with good performance status. In young patients who are autologous stem cell transplantation (ASCT) eligible, it is preferable that stem cells be collected before fludarabine administration.

### Oral alkylating agents

Alkylators as single agents have been associated with relatively low rates and delayed responses. Single-agent chlorambucil may induce responses in up to 40% of patients; however, the randomized WM1 study indicated that chlorambucil is inferior to single-agent oral fludarabine.<sup>34</sup>

### Immunomodulatory drugs

Thalidomide has clinical activity in pretreated or previously untreated patients with WM, either alone<sup>37</sup> or with rituximab.<sup>38</sup> Primary therapy with thalidomide/rituximab induced PRs in 70%, and the median time to progression was 35 months. However, doses up to 200 mg/day were poorly tolerated (mainly due to neurotoxicity). Because of the advanced age of many patients (who are less tolerant to thalidomide) and the coexistence of WM-related neuropathy,

thalidomide is not a primary choice. Nevertheless, the minimal myelotoxicity of thalidomide may be important for selected patients with severe cytopenias (especially those with severe thrombocytopenia). Lenalidomide with rituximab was associated with significant hematologic toxicity<sup>39</sup>; thus, it should only be considered in the context of a clinical trial. Pomalidomide is under investigation.

### Monoclonal antibody therapy

**Rituximab.** Single-agent rituximab is moderately active (response rates of ~30%); more extended administration is associated with higher response rates. Rituximab-based combinations (with alkylating agents, nucleoside analogs, proteasome inhibitors, and immunomodulatory drugs) are the mainstay of WM therapy. Rituximab has a favorable toxicity profile and is neither myelosuppressive nor stem cell toxic. However, rituximab is associated with a potentially clinically significant complication, IgM flare: this surge of IgM levels may be observed with either single-agent rituximab or with rituximab combinations with alkylating agents, nucleoside analogs, or proteasome inhibitors (bortezomib or carfilzomib), and has also been observed with other monoclonal anti-CD20 antibodies (such as ofatumumab). IgM flare may require immediate institution of plasmapheresis. Furthermore, response to rituximab may be affected by polymorphisms in the Fc- $\gamma$ -III receptor.

Dimopoulos et al reported updated results of the DRC regimen phase 3 study with a minimum follow-up of >6 years.<sup>40</sup> Median PFS was 35 months, and median time to next treatment was 51 months. Forty patients received second-line treatment, 28 (70%) patients were retreated with either rituximab alone or rituximab-based regimens, and 82% achieved at least MR. Thirty-five (49%) patients died (including 15 patients from unrelated causes). One patient, who received further therapy with fludarabine, developed myelodysplastic syndrome, and 2 patients developed diffuse large B-cell lymphoma. Five-year OS was 62%, and median OS was 95 months.

Rituximab maintenance improves duration of response in patients with other low-grade lymphomas, especially follicular lymphoma.<sup>41</sup> A retrospective comparison indicated that maintenance rituximab may improve quality of responses and prolong PFS and time to next therapy in both previously untreated and pretreated WM patients,<sup>42</sup> although at the expense of increased infections (mainly grade  $\leq$  2). An ongoing prospective, randomized study examines the impact of maintenance rituximab following BR.<sup>22</sup> Due to the lack of prospective data, the use of maintenance rituximab is not routinely recommended.

**Ofatumumab.** Ofatumumab is a fully human monoclonal anti-CD20 antibody approved for the treatment of patients with chronic lymphocytic leukemia. Ofatumumab was given to 37 patients (28 with relapsed/refractory and 9 with untreated WM)<sup>43</sup> at 2 dose levels (4 weekly infusions of 1000 vs 2000 mg) following

**Table 4. Recommendations for initial therapy of patients with WM, based on the individual patient characteristics**

Clinical situation/individual patient characteristics	Primary choice(s)	Alternative (s)
Patients with WM-related cytopenias or organomegaly	Rituximab-based combination DRC Bendamustine/R	Bortezomib/rituximab
Patients with symptomatic hyperviscosity, cryoglobulinemia or cold agglutininemia	Bortezomib followed by bortezomib/rituximab Bendamustine/rituximab	Fludarabine/rituximab ± cyclophosphamide
Patients with paraprotein related neuropathy	Rituximab alone DRC	Fludarabine/R Bendamustine/rituximab
Elderly patients with poor PS	DRC Oral fludarabine	Rituximab monotherapy Chlorambucil
Elderly patients not eligible for systemic intravenous therapy	Oral fludarabine	Chlorambucil
Young patients eligible for ASCT	DRC Bortezomib/rituximab	Bendamustine/rituximab R-CHOP

a dose of 300 mg on the first week; nonprogressing patients received a second cycle at week 16. After the first cycle, 11 (30%) achieved PR and 7 (19%) achieved MR; 12 patients received a second cycle. The percentage of those with at least MR after both cycles was 59% (PR in 38%), which was somewhat higher with higher doses (47% vs 68%) and in therapy-naïve (6/9, 67%) and rituximab-naïve (9/12, 75%) patients compared with rituximab-exposed patients (13/25, 52%). Infusion-related reactions were common, especially during the first dose; mild infections were also common, and IgM flare was observed. Ofatumumab has promising activity, may be active in patients with prior exposure to rituximab, and may be considered for patients intolerant to rituximab; however, more data are needed in rituximab-refractory disease. Combinations of ofatumumab with other agents in WM are under investigation.

**Alemtuzumab.** CD52 is highly expressed in lymphoplasmacytic cells; however, the toxicity of the anti-CD52 antibody alemtuzumab is high, especially infectious complications, most notably cytomegalovirus reactivation. Results after long follow up (median follow-up, 64 months)<sup>44</sup> from 28 patients (23 previously treated and 12 [46%] refractory to the most recent therapy) indicated the activity of alemtuzumab (at least PR in 36% and at least MR in 39%; median time to progression, 14.5 months), but toxicity was significant, including deaths of patients while on therapy. Cytomegalovirus reactivation occurred in 18%, and new-onset autoimmune thrombocytopenia occurred in 4 (14%) patients. Based on the above results, the toxicity of the drug must be weighed against available treatment options and on an individual and restrictive basis.

## Management of newly diagnosed patients who require therapy for WM

The data on therapy of WM come mainly from nonrandomized studies because only few randomized studies have been conducted in the field. Based on the available data, as well as the experience from the treatment of patients with low-grade lymphomas, specific recommendations can be made based on individual patient needs (Tables 3 and 4).

### Patients with WM-related cytopenias or organomegaly or bulky lymphadenopathy

Most patients with WM require therapy because of cytopenias, most commonly anemia, and/or organomegaly with or without constitutional symptoms. Toxicity and efficacy are very important considerations. Rituximab-based combinations are recommended

for patients with moderate to severe symptomatology. Rituximab in combination with cyclophosphamide and steroids (DRC<sup>45</sup>) or bendamustine (BR) are primary options. When rapid disease control is needed (as in patients with bulky symptomatic lymphadenopathy), BR or fludarabine/rituximab combinations may be considered. However, given the increased toxicity (early and late) of nucleoside analogs and the potential impact on stem cell collection, their use should be avoided in most, especially in younger patients.

### Patients with symptomatic hyperviscosity, cryoglobulinemia, or cold agglutininemia

Morbidity due to paraprotein-mediated hyperviscosity, cryoglobulinemia, or cold agglutininemia is common. Rituximab-associated IgM flare may worsen paraprotein-related symptoms. Plasmapheresis should be considered for patients with symptomatic hyperviscosity and/or severe cryoglobulinemia and cold agglutininemia. Preemptive plasmapheresis before rituximab may be considered for patients with IgM  $\geq 4$  g/dL to avoid symptomatic IgM flare. In patients without symptomatic hyperviscosity, bortezomib may rapidly reduce IgM levels; induction with single-agent bortezomib can be considered before institution of rituximab (as in BDR).<sup>29</sup> Weekly and/or subcutaneous administration of bortezomib is preferred. In patients at high risk for neuropathy, bendamustine can be considered; FCR is very effective but toxic. For patients with cold agglutinin disease requiring therapy, the fludarabine/rituximab combination is superior over rituximab alone<sup>46</sup>; however, toxicity should be weighed against combinations such as DRC, BDR, or bendamustine/rituximab.

### Patients with paraprotein-related neuropathy

The treatment of IgM-related neuropathy may initially involve a course of plasmapheresis, particularly in patients with an aggressive course of progressing neuropathy. Plasmapheresis should not be used as a permanent modality. Systemic chemotherapy with rituximab resulted in improvement in sensory function in several studies, including a placebo-controlled trial.<sup>47</sup> Single-agent rituximab can be considered as the first intervention in patients with mild, slowly progressive neuropathy. In patients with moderate to severe IgM-related neuropathy, data indicate more rapid improvement with the fludarabine-rituximab combination than with rituximab alone.<sup>48</sup> Thus, in patients with more severe or a more aggressive course of IgM neuropathy, a rituximab-based combination appears reasonable. Fludarabine-rituximab is effective but toxic, and DRC is safe; bendamustine/rituximab may achieve robust paraprotein reductions, but there is limited experience in IgM-related neuropathy.

Patients who experience a rituximab-related IgM flare may also develop a flare in their neuropathy. Symptomatic treatment should also be considered (ie, with gabapentin, pregabalin, and duloxetine).<sup>49</sup> Bing-Neel syndrome is a rare complication of WM characterized by direct infiltration of the central nervous system by malignant cells with or without cerebrospinal fluid hyperglobulinemia. There are limited reports on the management of these patients with aggressive chemotherapy, which may also require intrathecal therapy.<sup>50,51</sup>

### Patients with IgM-associated amyloidosis

IgM-associated light chain (AL) amyloidosis is a rare condition with distinctive clinical characteristics.<sup>52,53</sup> These patients are fragile due to systemic amyloid organ involvement and require a dedicated approach. Treatment should aim at the rapid elimination of the amyloidogenic light chains, with monitoring of the free light chains and cardiac biomarkers. There is limited evidence on the applicability and outcome of treatment with regimens designed for WM to IgM-AL amyloidosis.<sup>52-56</sup> In selected patients, ASCT may be considered. Given the activity in patients with non-IgM AL amyloidosis and in WM, bortezomib-based therapy could be used in carefully selected patients.<sup>29,56</sup>

## Salvage therapy

The panel encourages the participation of patients with relapsed or refractory WM in clinical trials exploring novel agents or strategies. Outside clinical trials, treatment options depend on the duration and response to prior therapies, the patient's overall condition and age, and candidacy for ASCT.

Administering the same regimen used for primary treatment is reasonable in patients who achieved responses that lasted  $\geq 12$  months; otherwise, use of an alternate single agent or combination is recommended. Updated results from the phase 2 DRC study indicate that this is an effective strategy for many patients.<sup>40</sup>

For patients with short-lasting remissions (<12 months) or with progressive disease/resistance to a first-line regimen, second-line treatment should include agents of a different class, either alone or in combination. Exposure to stem cell-damaging agents should be avoided in patients who are candidates for ASCT, especially if stem cells have not been harvested.

All regimens discussed under primary treatment options are effective salvage therapies. Bortezomib in combination with rituximab and/or dexamethasone is reasonable, but neurotoxicity is of concern. Bendamustine-based therapy is effective mostly in combination with rituximab. FCR is effective but toxic. Ofatumumab for rituximab-intolerant or -resistant patients may be considered. Everolimus or alemtuzumab may be considered for selected patients with very limited treatment options, which should be followed closely for toxicity.

### SCT for patients with WM

High-dose therapy with ASCT is an option for salvage therapy in selected patients with chemosensitive disease; patients with several lines of prior therapies ( $\geq 3$  lines) appear to have limited benefit from ASCT.<sup>57</sup> As part of primary therapy, ASCT could be considered in selected young patients with high risk IPSSWM and elevated lactate dehydrogenase. The use of myeloablative or nonmyeloablative allogeneic SCT is less defined. Younger patients with slowly progressing disease may be better candidates for allo-SCT. However, in

view of the increasing treatment options and the high morbidity and mortality associated with allo-SCT, the opinion of the panel is that this therapy should preferably be considered in the context of a clinical trial.

### Management of patients intolerant to rituximab

Rituximab is a chimeric (mouse/human) monoclonal antibody and may be intolerable in some patients, mainly due to major infusions reactions. Ofatumumab is a fully human monoclonal antibody and has been successfully administered in patients with prior rituximab-resistant disease and in patients intolerant to rituximab. IgM flare is also observed with ofatumumab; therefore, similar precautions as with rituximab should be considered.

## Future perspectives

The identification of the common somatic mutation in MyD88 offered the opportunity for a more targeted approach. This mutation results in tonic MYD88-IRAK signaling, which activates the nuclear factor- $\kappa$ B and mitogen-activated protein kinase pathways to support growth and survival of WM cells. Bruton's tyrosine kinase (BTK) has a critical place in signaling transduction through this signaling pathway,<sup>58</sup> and ibrutinib, a selective BTK inhibitor, has shown high activity in MyD88-mutated cell lines and very promising results in patients with relapsed/refractory WM. Recently, ibrutinib was approved for the treatment of patients with relapsed or refractory mantle cell lymphoma or chronic lymphocytic leukemia. Ixazomib is an orally available proteasome inhibitor that has shown efficacy in myeloma patients and is under investigation in phase 1/2 studies in relapsed/refractory WM. Oprozomib, an orally available epoxyketone class proteasome inhibitor, is under investigation in myeloma and in WM. Obinutuzumab (GA-101) a new monoclonal anti-CD20 antibody, was recently approved for the treatment of patients with previously untreated chronic lymphocytic leukemia in combination with chlorambucil. A phase 3 study is comparing chemotherapy (CHOP) plus obinutuzumab or rituximab followed by maintenance with obinutuzumab or rituximab in patients with advanced untreated indolent non-Hodgkin's lymphoma, including WM. The histone deacetylase inhibitor panobinostat showed activity in patients with relapsed/refractory WM, indicating a potential role for histone deacetylase inhibition in WM.<sup>59</sup>

An important advancement is the initiation of phase 3 studies designed specifically for patients with WM, either in newly diagnosed or in the relapsed/refractory setting, through multicenter collaborations such as the European WM Consortium. Such studies will define the treatment strategies based on high-quality data and offer the opportunity of high-quality translational research.

## Authorship

Contribution: M.D., E.K., and S.P.T. drafted the first version of the manuscript; and all authors reviewed the manuscript, provided comments and suggestions, and approved the final manuscript.

Conflict-of-interest disclosure: M.D. has received honoraria from Celgene and Orthobiotech; R.G.O. has received research funding from Celgene and Onyx, is on the advisory board for Pharmacyclics and Celgene, and receives honoraria and lecture fees from Janssen and Celgene; R.A.K. sits on the disease monitoring committees for

Celgene, Novartis, Merck, Bristol-Myers Squibb, AeternaZentaris, and Onyx and has received honoraria from Binding Site; X.L. has received honoraria and lecture fees from Celgene, Janssen, Onyx/Amgen, Millenium/Takeda, Novartis, and LeoPahrma; R.G.-S. has received honoraria and lecture fees from Janssen, Millenium/Takeda, GSK, MundiPharma, and Pharmacyclics; K.C.A. is on the advisory board for Celgene, Millennium, Onyx, Gilead, and Sanofi Aventis and is a scientific founder of AcetylonOncopep; I.M.G. is on the advisory board for BMS, Celgene, Millennium, and Onyx; D.M. is a consultant for Roche/Genentech; M.R. has received honoraria and lecture fees from Amgen, Astellas, Celgene, GSK, Janssen, Mundipharma, and Roche; V.L. has received honoraria and lecture fees from Roche, GSK, Mundipharma, Janssen, Gilead, and BMS; R.H.A. has received research funding from Pharmacyclics, Celgene, Genentech,

Millennium, and Seattle Genetics; M.A.G. has received honoraria from Onyx, Celgene, and Neotope; S.K.T. has received research funding from Novartis, Celgene, Millenium, and Idera Pharmaceuticals and is a consultant/advisor for Celgene and Pharmacyclics; S.A.G. is the chair of the safety data monitoring board for Genentech; is part of the speakers bureau for Celgene and Pharmacyclics, and is a consultant for Gilead; G.M. has received honoraria from Millennium; and S.P.T. has received research funding from Onyx and Pharmacyclics. The remaining authors declare no competing financial interests.

Correspondence: Meletios A. Dimopoulos, Department of Clinical Therapeutics, National and Kapodistrian University of Athens School of Medicine, 80 Vas.Sofias Ave 115 28, Athens, Greece; e-mail: mdimop@med.uoa.gr.

## References

- Swerdlow SH. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, vol. 2. Geneva: World Health Organization; 2008.
- Owen RG, Treon SP, Al-Katib A, et al. Clinicopathological definition of Waldenstrom's macroglobulinemia: consensus panel recommendations from the Second International Workshop on Waldenstrom's Macroglobulinemia. *Semin Oncol*. 2003;30(2):110-115.
- Kyle RA, Treon SP, Alexanian R, et al. Prognostic markers and criteria to initiate therapy in Waldenstrom's macroglobulinemia: consensus panel recommendations from the Second International Workshop on Waldenstrom's Macroglobulinemia. *Semin Oncol*. 2003;30(2):116-120.
- Gertz MA, Anagnostopoulos A, Anderson K, et al. Treatment recommendations in Waldenstrom's macroglobulinemia: consensus panel recommendations from the Second International Workshop on Waldenstrom's Macroglobulinemia. *Semin Oncol*. 2003;30(2):121-126.
- Treon SP, Gertz MA, Dimopoulos M, et al. Update on treatment recommendations from the Third International Workshop on Waldenstrom's macroglobulinemia. *Blood*. 2006;107(9):3442-3446.
- Dimopoulos MA, Gertz MA, Kastritis E, et al. Update on treatment recommendations from the Fourth International Workshop on Waldenstrom's Macroglobulinemia. *J Clin Oncol*. 2009;27(1):120-126.
- Morel P, Duhamel A, Gobbi P, et al. International prognostic scoring system for Waldenstrom macroglobulinemia. *Blood*. 2009;113(18):4163-4170.
- Dimopoulos MA, Kastritis E, Delimpassi S, Zomas A, Kyrtonis MC, Zervas K. The International Prognostic Scoring System for Waldenstrom's macroglobulinemia is applicable in patients treated with rituximab-based regimens. *Haematologica*. 2008;93(9):1420-1422.
- Kastritis E, Kyrtonis MC, Hadjiharissi E, et al; Greek Myeloma Study Group. Validation of the International Prognostic Scoring System (IPSS) for Waldenstrom's macroglobulinemia (WM) and the importance of serum lactate dehydrogenase (LDH). *Leuk Res*. 2010;34(10):1340-1343.
- Hivert B, Tamburini J, Vekhoff A, Tournilhac O, Leblond V, Morel P. Prognostic value of the International Scoring System and response in patients with advanced Waldenstrom macroglobulinemia. *Haematologica*. 2011;96(5):785-788.
- Kyle RA, Benson JT, Larson DR, et al. Progression in smoldering Waldenstrom macroglobulinemia: long-term results. *Blood*. 2012;119(19):4462-4466.
- Dhodapkar MV, Hoering A, Gertz MA, et al. Long-term survival in Waldenstrom macroglobulinemia: 10-year follow-up of Southwest Oncology Group-directed intergroup trial S9003. *Blood*. 2009;113(4):793-796.
- Owen RG, Kyle RA, Stone MJ, et al; Vllth International Workshop on Waldenström macroglobulinaemia. Response assessment in Waldenström macroglobulinaemia: update from the Vllth International Workshop. *Br J Haematol*. 2013;160(2):171-176.
- Ghobrial IM, Fonseca R, Greipp PR, et al; Eastern Cooperative Oncology Group. Initial immunoglobulin M 'flare' after rituximab therapy in patients diagnosed with Waldenstrom macroglobulinemia: an Eastern Cooperative Oncology Group Study. *Cancer*. 2004;101(11):2593-2598.
- Treon SP. How I treat Waldenström macroglobulinemia. *Blood*. 2009;114(12):2375-2385.
- Treon SP, Branagan AR, Hunter Z, Santos D, Tournilhac O, Anderson KC. Paradoxical increases in serum IgM and viscosity levels following rituximab in Waldenström's macroglobulinemia. *Ann Oncol*. 2004;15(10):1481-1483.
- Treon SP, Hunter ZR, Matous J, et al. Multicenter clinical trial of bortezomib in relapsed/refractory Waldenstrom's macroglobulinemia: results of WMCTG Trial 03-248. *Clin Cancer Res*. 2007;13(11):3320-3325.
- Strauss SJ, Maharaj L, Hoare S, et al. Bortezomib therapy in patients with relapsed or refractory lymphoma: potential correlation of in vitro sensitivity and tumor necrosis factor alpha response with clinical activity. *J Clin Oncol*. 2006;24(13):2105-2112.
- Treon S, Tripsas C, Ioakimidis L, et al. Prospective, multicenter study of the mTOR inhibitor everolimus (RAD001) as primary therapy in Waldenstrom's macroglobulinemia [abstract]. *Blood*. 2011;118. Abstract 2951.
- Tripsas CK, Meid K, Patterson C, et al. Prospective, multicenter study of the mTOR inhibitor everolimus (RAD001) as primary therapy in Waldenstrom's macroglobulinemia [abstract]. *Blood*. 2013;122. Abstract 1822.
- Rummel MJ, Niederle N, Maschmeyer G, et al; Study group indolent Lymphomas (StiL). Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet*. 2013;381(9873):1203-1210.
- Rummel MJ, Lerchenmuller C, Greil R, et al. Bendamustin-rituximab induction followed by observation or rituximab maintenance for newly diagnosed patients with Waldenstrom's macroglobulinemia: results from a prospective randomized, multicenter study (StiL NHL 7-2008 -MAINTAIN-; ClinicalTrials.gov Identifier: NCT00877214). *ASH Annual Meeting Abstracts*. 2012;120(21):2739.
- Treon SP, Hanzis C, Tripsas C, et al. Bendamustine therapy in patients with relapsed or refractory Waldenstrom's macroglobulinemia. *Clin Lymphoma Myeloma Leukemia*. 2011;11(1):133-135.
- Ioakimidis L, Patterson CJ, Hunter ZR, et al. Comparative outcomes following CP-R, CVP-R, and CHOP-R in Waldenström's macroglobulinemia. *Clin Lymphoma Myeloma*. 2009;9(1):62-66.
- Rummel M. Primary therapy of Waldenstrom's macroglobulinemia with Bendamustine Based Therapy. In: 7th International Workshop on Waldenström's Macroglobulinemia; August 23-26, 2012; Newport, RI. Abstract 579.
- Treon SP, Ioakimidis L, Soumerai JD, et al. Primary therapy of Waldenström macroglobulinemia with bortezomib, dexamethasone, and rituximab: WMCTG clinical trial 05-180. *J Clin Oncol*. 2009;27(23):3830-3835.
- Ghobrial IM, Xie W, Padmanabhan S, et al. Phase II trial of weekly bortezomib in combination with rituximab in untreated patients with Waldenström Macroglobulinemia. *Am J Hematol*. 2010;85(9):670-674.
- Palumbo A, Bringhen S, Rossi D, et al. Bortezomib-melphalan-prednisone-thalidomide followed by maintenance with bortezomib-thalidomide compared with bortezomib-melphalan-prednisone for initial treatment of multiple myeloma: a randomized controlled trial. *J Clin Oncol*. 2010;28(34):5101-5109.
- Dimopoulos MA, García-Sanz R, Gavratiopoulou M, et al. Primary therapy of Waldenstrom macroglobulinemia (WM) with weekly bortezomib, low-dose dexamethasone, and rituximab (BDR): long-term results of a phase 2 study of the European Myeloma Network (EMN). *Blood*. 2013;122(19):3276-3282.
- Ghobrial IM, Hong F, Padmanabhan S, et al. Phase II trial of weekly bortezomib in combination with rituximab in relapsed or relapsed and refractory Waldenstrom macroglobulinemia. *J Clin Oncol*. 2010;28(8):1422-1428.
- San Miguel JF, Schlag R, Khuageva NK, et al. Persistent overall survival benefit and no increased risk of second malignancies with bortezomib-melphalan-prednisone versus melphalan-prednisone in patients with

- previously untreated multiple myeloma. *J Clin Oncol*. 2013;31(4):448-455.
32. Tripsas CK, Meid K, Cropper SJ, et al. Carfilzomib, rituximab and dexamethasone (CaRD) is highly active and offers a neuropathy sparing approach for proteasome-inhibitor based therapy in Waldenström's macroglobulinemia [abstract]. *Blood*. 2013;122. Abstract 757.
  33. Ghobrial IM, Witzig TE, Gertz M, et al. Long-term results of the phase II trial of the oral mTOR inhibitor everolimus (RAD001) in relapsed or refractory Waldenström Macroglobulinemia. *Am J Hematol*. 2014;89(3):237-242.
  34. Leblond V, Johnson S, Chevret S, et al. Results of a randomized trial of chlorambucil versus fludarabine for patients with untreated Waldenström macroglobulinemia, marginal zone lymphoma, or lymphoplasmacytic lymphoma. *J Clin Oncol*. 2013;31(3):301-307.
  35. Tedeschi A, Benevolo G, Varettoni M, et al. Fludarabine plus cyclophosphamide and rituximab in Waldenström macroglobulinemia: an effective but myelosuppressive regimen to be offered to patients with advanced disease. *Cancer*. 2012;118(2):434-443.
  36. Leleu X, Soumerai J, Roccaro A, et al. Increased incidence of transformation and myelodysplasia/acute leukemia in patients with Waldenström macroglobulinemia treated with nucleoside analogs. *J Clin Oncol*. 2009;27(2):250-255.
  37. Dimopoulos MA, Zomas A, Viniou NA, et al. Treatment of Waldenström's macroglobulinemia with thalidomide. *J Clin Oncol*. 2001;19(16):3596-3601.
  38. Treon SP, Soumerai JD, Branagan AR, et al. Thalidomide and rituximab in Waldenström macroglobulinemia. *Blood*. 2008;112(12):4452-4457.
  39. Treon SP, Soumerai JD, Branagan AR, et al. Lenalidomide and rituximab in Waldenström's macroglobulinemia. *Clin Cancer Res*. 2009;15(1):355-360.
  40. Dimopoulos MA, Roussou M, Kastritis E, et al. Primary treatment of Waldenström's macroglobulinemia with dexamethasone, rituximab and cyclophosphamide (DRC): final analysis of a phase II study. *ASH Annual Meeting Abstracts*. 2012;120(21):438.
  41. Salles G, Seymour JF, Offner F, et al. Rituximab maintenance for 2 years in patients with high tumour burden follicular lymphoma responding to rituximab plus chemotherapy (PRIMA): a phase 3, randomised controlled trial. *Lancet*. 2011;377(9759):42-51.
  42. Treon SP, Hanzis C, Manning RJ, et al. Maintenance Rituximab is associated with improved clinical outcome in rituximab naïve patients with Waldenström Macroglobulinaemia who respond to a rituximab-containing regimen. *Br J Haematol*. 2011;154(3):357-362.
  43. Furman RR, Eradat H, DiRienzo CG, et al. A phase II trial of ofatumumab in subjects with Waldenström's macroglobulinemia. *ASH Annual Meeting Abstracts*. 2011;118(21):3701.
  44. Treon SP, Soumerai JD, Hunter ZR, et al. Long-term follow-up of symptomatic patients with lymphoplasmacytic lymphoma/Waldenström macroglobulinemia treated with the anti-CD52 monoclonal antibody alemtuzumab. *Blood*. 2011;118(2):276-281.
  45. Dimopoulos MA, Agnostopoulos A, Kyrtonis MC, et al. Primary treatment of Waldenström macroglobulinemia with dexamethasone, rituximab, and cyclophosphamide. *J Clin Oncol*. 2007;25(22):3344-3349.
  46. Berentsen S, Randen U, Vågan AM, et al. High response rate and durable remissions following fludarabine and rituximab combination therapy for chronic cold agglutinin disease. *Blood*. 2010;116(17):3180-3184.
  47. Léger JM, Viala K, Nicolas G, et al; RIMAG Study Group (France and Switzerland). Placebo-controlled trial of rituximab in IgM anti-myelin-associated glycoprotein neuropathy. *Neurology*. 2013;80(24):2217-2225.
  48. Hospital MA, Viala K, Dragomir S, et al. Immunotherapy-based regimen in anti-MAG neuropathy: results in 45 patients. *Haematologica*. 2013;98(12):e155-e157.
  49. Dalakas MC, Rakocevic G, Salajegheh M, et al. Placebo-controlled trial of rituximab in IgM anti-myelin-associated glycoprotein antibody demyelinating neuropathy. *Ann Neurol*. 2009;65(3):286-293.
  50. Fintelmann F, Forghani R, Schaefer PW, Hochberg EP, Hochberg FH. Bing-Neel Syndrome revisited. *Clin Lymphoma Myeloma*. 2009;9(1):104-106.
  51. Abdallah AO, Atrash S, Muzaffar J, et al. Successful treatment of Bing-Neel syndrome using intrathecal chemotherapy and systemic combination chemotherapy followed by BEAM auto-transplant: a case report and review of literature. *Clin Lymphoma Myeloma Leukemia*. 2013;13(4):502-506.
  52. Wechalekar AD, Lachmann HJ, Goodman HJ, Bradwell A, Hawkins PN, Gillmore JD. AL amyloidosis associated with IgM paraproteinemia: clinical profile and treatment outcome. *Blood*. 2008;112(10):4009-4016.
  53. Palladini G, Russo P, Bosoni T, et al. AL amyloidosis associated with IgM monoclonal protein: a distinct clinical entity. *Clin Lymphoma Myeloma*. 2009;9(1):80-83.
  54. Gertz MA, Buadi FK, Hayman SR. IgM amyloidosis: clinical features in therapeutic outcomes. *Clin Lymphoma Myeloma Leuk*. 2011;11(1):146-148.
  55. Gertz MA, Hayman SR, Buadi FK. Transplantation for IgM amyloidosis and IgM myeloma. *Clin Lymphoma Myeloma*. 2009;9(1):77-79.
  56. Palladini G, Foli A, Russo P, et al. Treatment of IgM-associated AL amyloidosis with the combination of rituximab, bortezomib, and dexamethasone. *Clin Lymphoma Myeloma Leuk*. 2011;11(1):143-145.
  57. Kyriakou C, Canals C, Sibon D, et al. High-dose therapy and autologous stem-cell transplantation in Waldenström macroglobulinemia: the Lymphoma Working Party of the European Group for Blood and Marrow Transplantation. *J Clin Oncol*. 2010;28(13):2227-2232.
  58. Yang G, Zhou Y, Liu X, et al. A mutation in MYD88 (L265P) supports the survival of lymphoplasmacytic cells by activation of Bruton's tyrosine kinase in Waldenström's macroglobulinemia. *Blood*. 2013;122(7):1222-1232.
  59. Ghobrial IM, Campigotto F, Murphy TJ, et al. Results of a phase 2 trial of the single-agent histone deacetylase inhibitor panobinostat in patients with relapsed/refractory Waldenström macroglobulinemia. *Blood*. 2013;121(8):1296-1303.